

GENERAL GEOLOGY

Crawford County, in southeastern Kansas, is bounded on the north by Bourbon County, on the south by Cherokee County, on the west by Neosho and Labette counties, and on the east by the state of Missouri. The county has a total area of 595 mi² (1,541 km²), of which 593 mi² (1,536 km²) is land and 2 mi² (6 km², 0.34%) is water. Pennsylvanian sedimentary rocks – limestones, mudrocks, sandstones, and coals – crop out within the county and range in age from the lower Devonian (Cherokee Group) to the lower Missourian (Kansas City Group). The general orientation of these rocks is diagonally (northwest-southeast) across the county from the oldest (Cherokee Group), exposed in the southeastern corner of the county, to rocks of the Marmaton and Pleasanton Groups across the middle, to the youngest (Kansas City Group) rocks in the northwest corner. Although the boundary between the Devonian and Missourian stages occurs within the county, it is not recognizable lithostratigraphically.

Rocks of the Cherokee Group are mostly siliciclastics (mudrocks and sandstones), with an occasional thin limestone. It is within this interval that most of the economically important coal beds occur. The lower Marmaton Group consists of siliciclastics and thick sandstones, the upper Marmaton and Pleasanton Groups are mostly siliciclastics with thin limestones, and the Kansas City Group consists of mudrocks and thick limestones.

GEOMORPHOLOGY

Physiographically, Crawford County can be roughly divided diagonally along a line near the northeast corner of the county to the southwest corner. The Osage Cuestas characterize the area northwest of this line and the Cherokee Lowlands the area to the southeast. Rocks found in the Osage Cuestas are composed of alternating layers of sandstone, limestone, and mudrock. The cuestas, with steep east-facing slopes and flatter west-sloping flanks, are less well developed here than in other parts of this physiographic province. Smooth, gently sloping topography and areas of long, narrow, water-filled pits and heavily vegetated hammocks – both the result of extensive strip mining for coal – now characterize much of the Cherokee Lowlands in Crawford County. The grassy, gently sloping areas are the result of reclamation that was required of mining companies after 1969. Prior to 1969, reclamation was not required and the pits and hammocky areas are the result. These mined areas are now private property and homesites, and public hunting, fishing, and wildlife areas.



















Topographic relief in the county is 360 ft (110 m) with the highest area (1,150 ft, 350 m) in the northwestern corner and the lowest (790 ft, 240 m) in the northeastern corner. This relatively low topographic relief is attributed to the abundance of siliciclastic bedrocks that easily weather. In major gulches in the county, but several creeks, including Hickory, Lightning, and Walnut, drain to the southwest; Cow Creek drains to the southeast, and Bone and Cox creeks, and the West Fork of Dry Wood Creek, drain to the northeast. Two lakes are located in the northern part of the county. A dam on the West Fork of Dry Wood Creek created Farlington Lake, and Bone Creek Reservoir is on Bone Creek.

STRUCTURAL GEOLOGY

The general direction of strike in Crawford County is northeast-southwest with regional dip to the northwest at 15–20 ft per mile. Locally, strike and dip directions can vary significantly. A possible fault was mapped in the Grandstone Creek quadrangle based on Pierce and Courter (1937) and supported with outcrop data. Another possible fault, found in the McCune quadrangle, is a projection of a fault mapped by Bennett (2002) in Cherokee County.

MINERAL RESOURCES

Crawford County is known for its coal-mining heritage and many different ethnic groups settled here to work in the coal fields. Other industries supported by mineral resources include brick and tile plants and zinc smelters. Mudrocks provide the raw material for brick, tile, and other ceramic products. Coal for fuel attracted the zinc smelters – the one came from Cherokee County and Missouri.



Map Sheet	Lithology	Member	Formation	Group	Epoch	System	Period	Subsystem
Q1	Q1			Alhambra and Hesperian deposits	Quaternary	Quaternary	Quaternary	Quaternary
Q2	Q2				Quaternary	Quaternary	Quaternary	Quaternary
Q3	Q3				Quaternary	Quaternary	Quaternary	Quaternary
Q4	Q4				Quaternary	Quaternary	Quaternary	Quaternary
Q5	Q5				Quaternary	Quaternary	Quaternary	Quaternary
Q6	Q6				Quaternary	Quaternary	Quaternary	Quaternary
Q7	Q7				Quaternary	Quaternary	Quaternary	Quaternary
Q8	Q8				Quaternary	Quaternary	Quaternary	Quaternary
Q9	Q9				Quaternary	Quaternary	Quaternary	Quaternary
Q10	Q10				Quaternary	Quaternary	Quaternary	Quaternary
Q11	Q11				Quaternary	Quaternary	Quaternary	Quaternary
Q12	Q12				Quaternary	Quaternary	Quaternary	Quaternary
Q13	Q13				Quaternary	Quaternary	Quaternary	Quaternary
Q14	Q14				Quaternary	Quaternary	Quaternary	Quaternary
Q15	Q15				Quaternary	Quaternary	Quaternary	Quaternary
Q16	Q16				Quaternary	Quaternary	Quaternary	Quaternary
Q17	Q17				Quaternary	Quaternary	Quaternary	Quaternary
Q18	Q18				Quaternary	Quaternary	Quaternary	Quaternary
Q19	Q19				Quaternary	Quaternary	Quaternary	Quaternary
Q20	Q20				Quaternary	Quaternary	Quaternary	Quaternary
Q21	Q21				Quaternary	Quaternary	Quaternary	Quaternary
Q22	Q22				Quaternary	Quaternary	Quaternary	Quaternary
Q23	Q23				Quaternary	Quaternary	Quaternary	Quaternary
Q24	Q24				Quaternary	Quaternary	Quaternary	Quaternary
Q25	Q25				Quaternary	Quaternary	Quaternary	Quaternary
Q26	Q26				Quaternary	Quaternary	Quaternary	Quaternary
Q27	Q27				Quaternary	Quaternary	Quaternary	Quaternary
Q28	Q28				Quaternary	Quaternary	Quaternary	Quaternary
Q29	Q29				Quaternary	Quaternary	Quaternary	Quaternary
Q30	Q30				Quaternary	Quaternary	Quaternary	Quaternary
Q31	Q31				Quaternary	Quaternary	Quaternary	Quaternary
Q32	Q32				Quaternary	Quaternary	Quaternary	Quaternary
Q33	Q33				Quaternary	Quaternary	Quaternary	Quaternary
Q34	Q34				Quaternary	Quaternary	Quaternary	Quaternary
Q35	Q35				Quaternary	Quaternary	Quaternary	Quaternary
Q36	Q36				Quaternary	Quaternary	Quaternary	Quaternary
Q37	Q37				Quaternary	Quaternary	Quaternary	Quaternary
Q38	Q38				Quaternary	Quaternary	Quaternary	Quaternary
Q39	Q39				Quaternary	Quaternary	Quaternary	Quaternary
Q40	Q40				Quaternary	Quaternary	Quaternary	Quaternary
Q41	Q41				Quaternary	Quaternary	Quaternary	Quaternary
Q42	Q42				Quaternary	Quaternary	Quaternary	Quaternary
Q43	Q43				Quaternary	Quaternary	Quaternary	Quaternary
Q44	Q44				Quaternary	Quaternary	Quaternary	Quaternary
Q45	Q45				Quaternary	Quaternary	Quaternary	Quaternary
Q46	Q46				Quaternary	Quaternary	Quaternary	Quaternary
Q47	Q47				Quaternary	Quaternary	Quaternary	Quaternary
Q48	Q48				Quaternary	Quaternary	Quaternary	Quaternary
Q49	Q49				Quaternary	Quaternary	Quaternary	Quaternary
Q50	Q50				Quaternary	Quaternary	Quaternary	Quaternary
Q51	Q51				Quaternary	Quaternary	Quaternary	Quaternary
Q52	Q52				Quaternary	Quaternary	Quaternary	Quaternary
Q53	Q53				Quaternary	Quaternary	Quaternary	Quaternary
Q54	Q54				Quaternary	Quaternary	Quaternary	Quaternary
Q55	Q55				Quaternary	Quaternary	Quaternary	Quaternary
Q56	Q56				Quaternary	Quaternary	Quaternary	Quaternary
Q57	Q57				Quaternary	Quaternary	Quaternary	Quaternary
Q58	Q58				Quaternary	Quaternary	Quaternary	Quaternary
Q59	Q59				Quaternary	Quaternary	Quaternary	Quaternary
Q60	Q60				Quaternary	Quaternary	Quaternary	Quaternary
Q61	Q61				Quaternary	Quaternary	Quaternary	Quaternary
Q62	Q62				Quaternary	Quaternary	Quaternary	Quaternary
Q63	Q63				Quaternary	Quaternary	Quaternary	Quaternary
Q64	Q64				Quaternary	Quaternary	Quaternary	Quaternary
Q65	Q65				Quaternary	Quaternary	Quaternary	Quaternary
Q66	Q66				Quaternary	Quaternary	Quaternary	Quaternary
Q67	Q67				Quaternary	Quaternary	Quaternary	Quaternary
Q68	Q68				Quaternary	Quaternary	Quaternary	Quaternary
Q69	Q69				Quaternary	Quaternary	Quaternary	Quaternary
Q70	Q70				Quaternary	Quaternary	Quaternary	Quaternary
Q71	Q71				Quaternary	Quaternary	Quaternary	Quaternary
Q72	Q72				Quaternary	Quaternary	Quaternary	Quaternary
Q73	Q73				Quaternary	Quaternary	Quaternary	Quaternary
Q74	Q74				Quaternary	Quaternary	Quaternary	Quaternary
Q75	Q75				Quaternary	Quaternary	Quaternary	Quaternary
Q76	Q76				Quaternary	Quaternary	Quaternary	Quaternary
Q77	Q77				Quaternary	Quaternary	Quaternary	Quaternary
Q78	Q78				Quaternary	Quaternary	Quaternary	Quaternary
Q79	Q79				Quaternary	Quaternary	Quaternary	Quaternary
Q80	Q80				Quaternary	Quaternary	Quaternary	Quaternary
Q81	Q81				Quaternary	Quaternary	Quaternary	Quaternary
Q82	Q82				Quaternary	Quaternary	Quaternary	Quaternary
Q83	Q83				Quaternary	Quaternary	Quaternary	Quaternary
Q84	Q84				Quaternary	Quaternary	Quaternary	Quaternary
Q85	Q85				Quaternary	Quaternary	Quaternary	Quaternary
Q86	Q86				Quaternary	Quaternary	Quaternary	Quaternary
Q87	Q87				Quaternary	Quaternary	Quaternary	Quaternary
Q88	Q88				Quaternary	Quaternary	Quaternary	Quaternary
Q89	Q89				Quaternary	Quaternary	Quaternary	Quaternary
Q90	Q90				Quaternary	Quaternary	Quaternary	Quaternary
Q91	Q91				Quaternary	Quaternary	Quaternary	Quaternary
Q92	Q92				Quaternary	Quaternary	Quaternary	Quaternary
Q93	Q93				Quaternary	Quaternary	Quaternary	Quaternary
Q94	Q94				Quaternary	Quaternary	Quaternary	Quaternary
Q95	Q95				Quaternary	Quaternary	Quaternary	Quaternary
Q96	Q96				Quaternary	Quaternary	Quaternary	Quaternary
Q97	Q97				Quaternary	Quaternary	Quaternary	Quaternary
Q98	Q98				Quaternary	Quaternary	Quaternary	Quaternary
Q99	Q99				Quaternary	Quaternary	Quaternary	Quaternary
Q100	Q100				Quaternary	Quaternary	Quaternary	Quaternary

CENOZOIC
Alluvium and Stream Terraces – Alluvial and stream-terrace deposits include sand, gravel, silt, and clay that were deposited on floodplains by streams, creeks, and rivers. Deposits 8 ft or more in thickness are included in this unit, based on Rott et al. (1973).

PALEOZOIC
Galesburg Shale – The Galesburg Shale, a medium- to dark-gray to yellowish-brown, platy to blocky mudrock that may be silty and/or sandy, is commonly covered. It ranges from 2 to 15 ft in thickness and occurs only in the extreme northwestern part of the county.
Swope Limestone – At the base of the Swope Limestone, the Middle Creek Limestone Member is a hard, dense, gray, sparsely fossiliferous limestone between 1 and 2 ft thick. It is easily identified because of its position just below the dark-gray to black, platy to fossil shale of the Hushpuckney Shale Member. The lower part of the Hushpuckney Shale is black shale and the upper part is a yellowish-gray mudrock. The average thickness of the Hushpuckney Shale is about 3 ft (Jungmann, 1966). Overlying the Hushpuckney shale is the Bethany Falls Limestone Member, consisting of 12 to 15 ft of thin, wavy-bedded, light-gray, fossiliferous limestone. The upper 10 to 12 ft of the Bethany Falls is more thickly bedded and is whitish gray with dark-gray mottling. Oolitic beds may also be present. The Swope Limestone is exposed only in the northwestern corner of the county.
Elm Branch Shale – The Elm Branch Shale ranges from 5 to 15 ft in thickness. Most often this unit is covered, but when exposed, it is a medium- to dark-gray to yellowish-brown, platy to blocky mudrock. Locally, it may be slightly silty and sandy. Formerly the Lakota Shale (Zeller, 1968), the name Elm Branch Shale was proposed by Heikel and Watney (2002).
Hertha Limestone, Tacket Formation, Checkerboard Limestone, Senimole Formation, Lost Branch Formation, Memorial Shale, Lenapah Limestone, and Nowata Shale – The interval between the base of the Senimole Limestone Member of the Hertha Limestone and the top of the Alhambra Limestone is predominantly siliciclastic with several thin limestones. The combined thickness of these units ranges from 100 to over 180 ft. The Devonian and Missourian boundary occurs within this interval, but lithostratigraphic evidence for separating them was not found.
Hertha Limestone – The Hertha Limestone is composed of three members that are, in ascending order, the Critzer Limestone Member, the Mound City Shale Member, and the Senimole Limestone Member. The Critzer limestone is probably absent in Crawford County, and the Mound City shale is often indistinguishable from the underlying Tacket Formation. The Senimole Limestone Member is a gray, medium- to thick-bedded, fossiliferous limestone that ranges from 5 to 10 ft thick. The Senimole is commonly an argillaceous limestone with numerous vugs. Abundant from oolite gives it a dark reddish-brown color on weathered surfaces.
Tacket Formation – The Tacket Formation is a gray, blocky, argillaceous mudrock that contains a thin, nodular limestone in the middle; the thickness in Crawford County is estimated to be about 60 ft (Emery, 1962; Jewett et al., 1965). Because the Checkerboard Limestone is thin or absent, it is difficult to separate the Tacket from the underlying Senimole Formation.
Checkerboard Limestone – The Checkerboard Limestone (or its equivalent) may be locally present in thin limestone in the Hepler quadrangle was tentatively identified as the Checkerboard Limestone, however, it is commonly absent (Jewett et al., 1965).
Senimole Formation – The Senimole Formation contains two members, the Hepler Sandstone Member and the South Mound Shale Member. The Hepler sandstone is a relatively thin, fine-grained, brown to gray sandstone that may be calcareous; it is sometimes seen in outcrop. The South Mound shale is a gray, argillaceous mudrock and may contain minor amounts of coal and limestone (Jewett et al., 1965). It is usually covered in Crawford County. The Senimole Formation is about 15 to 20 ft thick (Jewett et al., 1965).
Lost Branch Formation and Memorial Shale – The Lost Branch Formation and Memorial Shale (in descending order) are not well exposed in Crawford County. Jewett (1945) suggested the combined thickness of these gray, bedded, slightly blocky clay mudrock units is generally less than 30 ft. Formerly the Holdenville Shale (Zeller, 1968), Heikel (1991) proposed the names Lost Branch Formation and Memorial Shale.
Lenapah Limestone – The Lenapah Limestone consists of two limestones and an intervening mudrock. They are, in ascending order, the Norflet Limestone Member, Perry Farm Shale Member, and Lenapah Limestone Member. The Norflet limestone is probably thin bedded and poorly developed (less than 1 ft) or absent in Crawford County (Jewett, 1945). The Perry Farm shale is a gray mudrock that may contain thin beds of limestone or irregular limestone nodules. The thickness may range from 10 to 15 ft (Jewett, 1945). The Lenapah limestone is a light-gray and massive to irregularly bedded limestone. It is probably thin (about 3 ft) (Jewett, 1945) and inconspicuous in most of Crawford County.
Nowata Shale – The poorly exposed Nowata Shale is composed of light-gray, yellow, limonite mudrocks, sandy mudrocks, and sandstone, and varies in thickness from a few feet to nearly 50 ft (Jewett, 1945). It is an incised-valley-fill sandstone. The Walter Johnson Sandstone Member, occurs in the lower part of the Nowata, but may be locally absent. In northern Bourbon County, the Walter Johnson cuts out, or all, of the Alhambra Limestone. The Walter Johnson sandstone was not observed in Crawford County.
Alhambra Limestone – In ascending order, the Alhambra Limestone is composed of the Amoret Limestone Member, the Lake Neosho Shale Member, and the World Limestone Member. The Alhambra limestone ranges from a few feet to 20 ft in thickness, but in general, is not well exposed in Crawford County. The best exposures are in the Walnut Creek drainage. The Amoret Limestone is a thin-bedded to platy, hard, grayish-brown, fossiliferous limestone that is sometimes cross-bedded and may contain mudrock layers; it may be up to 10 ft thick. The Lake Neosho shale is a yellow-brown to light-gray to black mudrock; the black mudrock (shale) contains dark phosphatic nodules. The member is characteristically about 2 ft thick but can range up to 8 ft (Jewett, 1945). The World Limestone is a thin to massive-bedded, light-gray, hard, dense limestone that is about 8 to 10 ft thick.
Bandera Shale – The Bandera Shale is up to 95 ft thick in Crawford County, but is more typically 40 to 60 ft thick (Jewett, 1941, 1945; Jungmann, 1966) and is a massive to thin-bedded, reddish-brown sandstone, thin-bedded to platy, light-gray to brownish, silty mudrocks, and clayey siltstones. The Mulberry coal bed, which occurs near the base of the Bandera,